

Assessment of technical knowledge on oral nutritional therapy between physicians and the nursing team of a private institution

Avaliação do conhecimento técnico sobre terapia nutricional oral entre médicos e equipe de enfermagem de uma instituição privada

Caroline Rosa Koerner^{1*}, Josiane de Paula Freitas¹, Thais Manfrinato Miola¹

ABSTRACT

Introduction: Malnutrition affects an average of 20-80% of cancer patients, leading to an impaired healing process, increased infection rates and hospital costs, and negative impacts on clinical results. Oral nutritional supplements (ONS) help to increase the patients caloric and protein supplies, promoting recovery of nutritional status. In order to have good adherence to the ONS, it is necessary that the multidisciplinary team assigned is aligned with regard to the information transferred to the patient. **Objectives:** This study aimed to assess the technical knowledge about oral nutritional therapy (ONT) among physicians and the nursing staff at a private institution. **Materials and Methods:** This was a descriptive cross-sectional study. The physicians and residents as well as the nursing staff who work in clinical patient care were invited to answer a semi-structured questionnaire to assess their technical knowledge regarding ONT. **Results:** This study evaluated 105 professionals. Most physicians and nursing staff stated that ONS is a food supplement. Although they reported not having had technical training on the subject, most of them felt capable of prescribing and providing guidance participants neglected symptoms and conditions relevant to the nutritional status of the patient, such as ascites, enteral nutritional therapy and home parenteral nutrition, concomitant chemotherapy/radiotherapy, and emesis. **Conclusion:** Information about ONS among medical and nursing professionals varies greatly. Interdisciplinary technical training is essential for the success of ONT as a tool for preventing and combating malnutrition.

Keywords: Malnutrition; Oncology service, Hospital; Nutrition therapy; Patient care team; Dietary supplements.

RESUMO

Introdução: A desnutrição afeta em média 20-80% dos pacientes com câncer, levando a um processo de cicatrização prejudicado, aumento das taxas de infecção e custos hospitalares, além de impactos negativos nos resultados clínicos. Os suplementos nutricionais orais (SNO) auxiliam no aumento do aporte calórico e proteico do paciente, promovendo a recuperação do estado nutricional. Para uma boa adesão ao SNO, é necessário que a equipe multiprofissional alocada esteja alinhada com as informações repassadas ao paciente. **Objetivos:** Este estudo teve como objetivo avaliar o conhecimento técnico sobre terapia nutricional oral (TNO) entre médicos e equipe de enfermagem de uma instituição privada. **Material e Métodos:** Trata-se de um estudo descritivo transversal. Os médicos e residentes, bem como a equipe de enfermagem que atuam na assistência clínica ao paciente, foram convidados a responder a um questionário semiestruturado para avaliar seu conhecimento técnico sobre TNO. **Resultados:** Este estudo avaliou 105 profissionais. A maioria dos médicos e enfermeiros afirmou que o ONS é um suplemento alimentar. Apesar de relatarem não ter tido capacitação técnica sobre o assunto, a maioria se sentiu capaz de prescrever e orientar os participantes sobre sintomas negligenciados e condições pertinentes ao estado nutricional do paciente, como ascite, terapia nutricional enteral e parenteral domiciliar, quimioterapia concomitante/radioterapia e vômitos. **Conclusão:** As informações sobre SNO entre profissionais médicos e de enfermagem variam muito. A formação técnica interdisciplinar é essencial para o sucesso da TNO como ferramenta de prevenção e combate à desnutrição.

Descritores: Desnutrição; Serviço de oncologia hospitalar; Terapia nutricional; Equipe de atendimento ao paciente; Suplementos dietéticos.

1. AC Camargo Cancer Center, Nutrição - São Paulo - São Paulo - Brazil

Financial support: The authors declare that no funds, grants, or other support were received during the preparation of this manuscript.

Conflicts of interest: The authors have no relevant financial or non-financial interests to disclose.

Correspondence author: Thais Manfrinato Miola.

E-mail: thais.miola@accamargo.org.br

Received on: November 29, 2022 | **Accepted on:** February 1, 2023 | **Published on:** March 2, 2023

DOI: <https://doi.org/10.5935/2526-8732.20230391>



INTRODUCTION

Cancer patients face challenges that go beyond the tumor, including malnutrition. Malnutrition can be defined as a state resulting from a deficit of nutrients that leads to changes in body composition, mental state, and function; can damage the clinical outcome; and has a multifactorial etiology.^[1] Cederholm et al. (2015)^[2] suggest malnutrition as a body mass index (BMI) $<18.5\text{kg/m}^2$ or $\text{BMI}<20\text{kg/m}^2$; for patients older than 70 years old. A $\text{BMI}<22\text{kg/m}^2$ associated with weight loss of $>10\%$ for an unspecified time or $>5\%$ in the last 3 months can also be used to diagnose a patient as being malnourished. However, the Global Leadership Initiative on Malnutrition suggests that the diagnosis be made considering five criteria, which are divided into phenotypic and etiological criteria. The phenotypic criteria include involuntary weight loss, low BMI, and reduced muscle mass; meanwhile, the etiological criteria include reduced food intake or nutrient absorption, the presence of acute or chronic inflammation, and the severity of the disease. To be diagnosed with malnutrition, the patient must meet at least one phenotypic criterion and one etiological criterion.^[3]

Unintentional weight loss in hospitalized patients used in isolation is already a negative prognostic indicator, and studies indicate that approximately 31-87% of individuals diagnosed with cancer show substantial loss of body weight, which is a marker for malnutrition.^[4,5] Unlike malnutrition in noncancer patients, cancer-associated malnutrition results in a negative energy balance and loss of muscle mass caused not only by decreased food intake but also by metabolic disorders such as insulin resistance, increased basal metabolism rate, and increased lipolysis and proteolysis, which are due to the presence of systemic and catabolic inflammation that aggravate weight loss. Considering these factors, nutritional support is a strong ally to reverse malnutrition in cancer patients.^[6]

Currently, it is estimated that, on average, the malnutrition rate in cancer patients varies from 20% to 80%; in addition, approximately 20% of cancer patients die due to malnutrition/cachexia.^[7]

The main complications related to malnutrition include a delay in the healing process; impaired absorption and binding of proteins, hepatic metabolism, and renal elimination of drugs and their metabolites; worsening of the immune response; increased rates of infections and postoperative complications; increased risk of developing pressure injury; increased length of hospital stay; increased risk of mortality; and a considerable increase in hospital costs. In addition to negatively impacting patient metabolism, involution of the nutritional status can also affect cancer treatment outcomes.^[8,9]

An epidemiological study carried out in Spain showed through the Global Subjective Assessment produced by the patient (ASG-PPP) that 96.7% of the patients were at nutritional risk and lacked early nutritional intervention. The authors observed that

the patients with the greatest weight loss were those with tumors in the esophagus, stomach, or larynx.^[10]

Nutritional support aims to significantly improve the response to treatment, contain weight loss and malnutrition, reduce the risk of complications, reduce morbidity and mortality, and provide a better quality of life for cancer patients. Among nutritional interventions, oral nutritional therapy (ONT) has been highlighted as its main objective is to increase the patient's dietary intake.^[11-13]

Oral nutritional supplements (ONS) are the main pillar of ONT. Available for oral consumption, these supplements can be found in ready-to-eat models (in tetra-pack packaging or in cans) or in powder form for reconstitution, and they are intended to complement food intake. ONS are indicated for patients who do not achieve their food intake requirements or for those who show weight loss; thus, their use helps the patients to recover their nutritional status.^[14] Supplements are considered as the first option, since they have physiological properties and are less invasive than other treatment methods; however, they should only be used as an aid for recovery or maintenance of nutritional status and not as an exclusive method of feeding or replacing meals.^[13-15]

There are several types of nutritional supplements, which are divided into standard and specialized formulas, according to their purpose and composition. Currently, the nutritional supplement industry has a wide variety of products, including normal to hypercaloric; normal to hyperprotein; free of sucrose or sugar; fortified with immunomodulatory nutrients; free or high in fat; with or without flavor; with or without dietary fiber; and even products aimed at certain audiences or those with specific clinical conditions such as cancer patients, those with kidney or liver failure, among others.^[15] These products, in turn, help with several processes, namely muscle formation and function, cholesterol lowering, intestinal function, amino acid synthesis, cell division, immune system function, red blood cell formation, and energy metabolism including protein, carbohydrate, and fat metabolism.^[16]

Several studies indicate that nutritional intervention should be an adjunct to cancer treatment. Moreover, a multidisciplinary approach allows better adherence to ONT, which is a key factor for successful treatment and recovery of patients.^[17]

A previous study has evaluated 13 randomized clinical trials of oral nutritional intervention (nutrition counseling, ONT, or both) in patients diagnosed with cancer at nutritional risk/malnourished during anticancer treatment or palliative care and compared oral nutritional intervention with usual care.^[18] The findings suggest that interventions can improve the quality of life of patients who are at nutritional risk or are malnourished. Furthermore, the nutritional intervention was the most effective for patients whose disease is curable or for those who have a long disease-free period after treatment.

According to Paccagnella et al. (2010)^[19] early dietary intervention in patients diagnosed with head and neck cancer who were undergoing chemotherapy or radiotherapy resulted in a greater tolerance to treatment, improvement in symptoms, and less hospitalization. The results of this study suggest that this intervention should be started before treatment and should be continued until the end of the treatment.

Additionally, a retrospective study that aimed to evaluate the impact of nutritional supplementation on the length of hospital stay, hospital costs, and readmission of the patient showed that the use of ONS reduced the length of hospital stay by an average of 2-3 days, reduced hospital costs by US\$4,734, and decreased the probability of these patients being readmitted within 30 days of discharge.^[20]

In practice, ONT requires multidisciplinary work, considering that each specialty is responsible for a different part of the process. For example, the Medicine Department (except for professional nutritionists) is often the area in charge of prescribing supplements, and nurses and their assistants are responsible for their delivery and administration. However, these specialties have different views on supplementation; supplements can be considered as a medicine or as a product that supplements nutrition. The different understandings about the purpose of the supplement, when passed on to the patient, can cause some confusion about its real benefit; moreover, a variable representation about such a relevant treatment has an impact on the importance of ONT, resulting in increased efforts to be made to prevent or treat malnutrition and to plan follow-up care.^[21]

It is essential that nutritional care is present from the early stages of the disease's natural history, and its integration in the therapeutic strategy in the battle against cancer is essential.^[13-22] Considering that the success of ONT as a treatment as well as its use for the prevention of weight loss and malnutrition depends on a good understanding of the health care team working together, the present study aimed to assess the knowledge and conduct of professionals working with this resource.

MATERIAL AND METHODS

Study design and Data survey

This was a descriptive cross-sectional study. Data collection was performed through the application of a semi-structured questionnaire (Appendix 1 and 2) to residents and physicians as well as the nursing staff who work at the A.C. Camargo Cancer Center. This was a cross-sectional descriptive study. Data collection was performed by applying a semi-structured questionnaire (Appendix 1 and 2) to residents and physicians, as well as to nursing staff working at the A.C. Camargo Cancer Center.

Included in the study were professionals who perform direct care in the area of clinical oncology, considering physicians and medical residents, and the nursing staff, such as nurses and nursing technicians. Exclusion criteria were physicians from other specialties, such as surgeons and radiologists, and nursing staff who work in the intensive care unit, diagnostic and imaging department and teaching and research department.

All participants completed the free and informed consent form prior to completing the questionnaires. The administration of the questionnaires was carried out between October and December 2020.

Semi-structured questionnaire

Physicians and nurses were invited to answer a semi-structured questionnaire that included questions regarding their profession and specialty, what ONS means to them, knowledge of the importance of ONT for the patient, whether this topic was addressed during their professional training, modalities and personal safety for prescription (physicians), conduct towards ONT, self-confidence to resolve possible doubts from patients, and open questions/criticisms and suggestions if the professional feels the need to report something that was not addressed.

Statistical analysis

Statistical analysis was performed using the software package IBM SPSS Statistic, version 22.0 (Chicago, IL, USA). Continuous variables were presented as the median or mean and standard deviation for non-normally and normally distributed data, respectively. Normality was tested using the Kolmogorov-Smirnov test. Ordinal or nominal variables were presented in absolute numbers and as a percentage of the total. Absolute and relative frequencies were calculated for categorical variables. For the analysis of categorical variables, the chi-squared test or Fisher's exact test was used, when appropriate. Comparison of the averages of continuous variables between the evaluation methods was performed using the Student's t-test or the nonparametric Mann-Whitney test for independent samples, as appropriate.

RESULTS

A total of 105 professionals were included in this study, including 34 physicians (32.4%) and 71 nursing staff (67.6%). Among the nursing staff, 35 were nursing technicians (33.3%), 35 were nurses (33.3%), and only 1 was a nursing assistant (1%). In the physician category, 11 were physicians (10.5%), and 23 were residents (21.9%).

Tables 1 and 2 describe the frequency of responses by the physicians and nursing staff, in the proper order. The analysis of common questions is presented in Table 3.

Table 1. Summary of the answers given by the nursing team.

Variable	Category	N (%)	
Do you know what ONS is?	Medication	0	
	Meal replacement	1 (1.4)	
	Meal complement	68 (95.8)	
	I do not know	2 (2.8)	
Was this topic addressed during your professional training?	Yes	30 (42.9)	
	No	34 (48.6)	
	I do not know	6 (8.6)	
Who is this supplement for?	Underweight patients	31 (43.7)	
	Patients with associated diseases	8 (11.3)	
	I do not know	7 (9.9)	
	Malnourished patients	40 (56.3)	
	All cancer patients	9 (12.7)	
What criteria would you use to classify a patient at nutritional risk?	Weight loss	61 (85.9)	
	Low food acceptance	62 (87.3)	
	Dysphagia	38 (53.5)	
	Diarrhea	33 (46.5)	
	Vomiting	30 (42.3)	
	Mucositis	35 (49.3)	
	Head and neck cancer	30 (42.3)	
	Ascites	12 (16.9)	
	Enteral/Parenteral home therapy	20 (28.2)	
	Chemotherapy concomitant with radiotherapy	19 (26.8)	
	What benefits do you believe the ONS brings to cancer patients?	Weight gain	56 (78.9)
		Better food intake	37 (52.1)
Better tolerance to treatment		46 (64.8)	
Lower chances of hospital readmission		19 (26.8)	
Do you notice an improvement in the patient's general condition and/or food acceptance while using the ONS?	None	0	
	Yes	48 (67.6)	
	No	5 (7.0)	
By whom is ONS prescribed?	I do not know	18 (25.4)	
	Doctor	11 (15.5)	
	Nutricionist	62 (87.3)	
	I do not know	1 (1.4)	
Who is the professional responsible for delivering the ONS in your department?	Nutritional Doctor	31 (43.7)	
	Nurse	18 (25.4)	
	Nursing technician	62 (87.3)	
	Nursing assistant	25 (35.2)	
When administering the ONS to the patient, do you guide him?	Yes	53 (79.1)	
	No	10 (14.9)	
	I do not know	4 (6.0)	
If not, why?	It is another professional's responsibility	5 (35.7)	
	I do not know how to assess the need for the supplement	4 (28.6)	
	The nutritional status of the patient is not relevant to his/her treatment	0	
What do you do if the patient refuses the ONS?	I advise him/her to talk to the doctor	4 (5.6)	
	I advise him/her to talk to the nutritionist	47 (66.2)	
	I communicate the doctor	6 (8.5)	
	I communicate the nutritionist	40 (56.3)	

Legend: ONS: Oral nutritional supplements.

Table 2. Summary of the answers given by the medical team

Variable	Category	N (%)
Do you know what ONS is?	Medication	0
	Meal replacement	1 (2,9)
	Meal complement	33 (95,8)
	I do not know	0
Was this topic addressed during your professional training?	Yes	9 (26,5)
	No	25 (73,5)
	I do not know	0
Do you prescribe or have you already prescribe an ONS?	Yes	18 (52,9)
	No	16 (47,1)
Based on which diagnosis do you prescribe the ONS?	Underweight patients	12 (35,3)
	Patients with associated diseases	1 (2,9)
	I do not know	7 (20,6)
	Malnourished patients	20 (58,8)
	All cancer patients	2 (5,9)
What criteria would you use to classify a patient at nutritional risk?	Weight loss	34 (100,0)
	Low food acceptance	31 (91,2)
	Dysphagia	23 (67,6)
	Diarrhea	17 (50,0)
	Vomiting	16 (47,1)
	Mucositis	22 (64,7)
	Head and neck cancer	21 (61,8)
	Ascites	10 (29,4)
	Enteral/Parenteral home therapy	13 (38,2)
	Chemotherapy concomitant with radiotherapy	16 (47,1)
	When prescribing the ONS to the patient, do you guide him/her?	Yes
No		7 (20,6)
I do not prescribe ONS		14 (41,2)
If not, why?	It is another professional's responsibility	9 (42,9)
	I don't know how to assess the need for the supplement	12 (57,1)
	The nutritional status of the patient is not relevant to his/her treatment	0
What benefits do you believe the ONS brings to cancer patients?	Weight gain	26 (76,5)
	Better food intake	12 (35,3)
	Better tolerance to treatment	28 (82,4)
	Lower chances of hospital readmission	26 (76,5)
	None	0
Do you notice an improvement in the patient's general condition and/or food acceptance while using the ONS?	Yes	27 (79,4)
	No	1 (2,9)
	I do not know	6 (17,6)
What do you do if the patient refuses the ONS?	I suspend	1 (2,9)
	I advise him/her to talk to the nutritionist	21 (61,8)
	I communicate the nutritionist	4 (11,8)
	I reinforce the importance of the ONS	(23,5)

Legend: ONS: Oral nutritional supplements.

Table 3 - Association of common responses between teams

Variable	Category	Doc. N (%)	Nur. N (%)	p-value
Do you know what ONS is?	Medication	0	0	1,000
	Meal replacement	1 (2,9)	1 (1,4)	
	Meal complement	33 (95,8)	68 (95,8)	
	I do not know	0	2 (2,8)	
Was this topic addressed during your professional training?	Yes	9 (26,5)	30 (42,9)	0,028
	No	25 (73,5)	34 (48,6)	
	I do not know	0	6 (8,6)	
Based on which diagnosis do you prescribe the ONS?	Underweight patients	12 (35,3)	31 (43,7)	0,546
Who is this supplement for?	Patients with associated diseases	1 (2,9)	8 (11,3)	0,266
	I do not know	7 (20,6)	7 (9,9)	0,140
	Malnutrition patients	20 (58,8)	40 (56,3)	0,976
	All cancer patients	2 (5,9)	9 (12,7)	0,497
What criteria would you use to classify a patient at nutritional risk?	Weight loss	34 (100,0)	61 (85,9)	0,028
	Low food acceptance	31 (91,2)	62 (87,3)	0,747
	Dysphagia	23 (67,6)	38 (53,5)	0,245
	Diarrhea	17 (50,0)	33 (46,5)	0,897
	Vomiting	16 (47,1)	30 (42,3)	0,799
	Mucositis	22 (64,7)	35 (49,3)	0,203
	Head and neck cancer	21 (61,8)	30 (42,3)	0,096
	Ascites	10 (29,4)	12 (16,9)	0,223
	Enteral/Parenteral home therapy	13 (38,2)	20 (28,2)	0,415
	Chemotherapy concomitant with radiotherapy	16 (47,1)	19 (26,8)	0,065
What benefits do you believe the ONS brings to cancer patients?	Weight gain	26 (76,5)	56 (78,9)	0,979
	Better food intake	12 (35,3)	37 (52,1)	0,159
	Better tolerance to treatment	28 (82,4)	46 (64,8)	0,106
	Lower chances of hospital readmission	26 (76,5)	19 (26,8)	< 0,001*
	None	0	0	1,000
Do you notice an improvement in the patient's general condition and/or food acceptance while using the ONS?	Yes	27 (79,4)	48 (67,6)	0,502
	No	1 (2,9)	5 (7,0)	
	I do not know	6 (17,6)	18 (25,4)	
When prescribing the ONS to the patient, do you guide him/her? When administering the ONS to the patient, do you guide him/her?	Yes	13 (38,2)	53 (79,1)	< 0,001*
	No	7 (20,6)	10 (14,9)	
	I do not know	-	4 (6,0)	
	I do not prescribe ONS	14 (41,2)	-	
If not, why?	It's another professional's assignment	9 (42,9)	5 (35,7)	0,012
	I don't know how to assess the need for the supplement	12 (57,1)	4 (28,6)	
	The nutritional status of the patient is not relevant to his treatment	0	0	

Legend: ONS, oral nutritional supplements; Doc, doctors; Nur, nursing;

*p-value < 0,005 According to Fisher's exact test.

DISCUSSION

The adherence of patients to ONT as a way to prevent and cope with malnutrition is directly linked to the knowledge of the team involved in the prescription, administration, and guidance of the patient, since they are responsible for transmitting all the necessary information for such a treatment. A team that does not have a unified view can create confusion for the main beneficiary of the treatment: the patient; this can lead to a reduced effectiveness and a poor adherence to the treatment by the patient.^[21]

A systematic review by Baldwin et al. (2012)^[18] suggests that the use of ONS is associated with significant reductions in complications associated with unintentional weight loss, length of hospital stay, and mortality rates. Therefore, nutritional assistance is a great ally against malnutrition. In addition, the report on recommending procedures for ONS for malnourished clinical or surgical patients or those at risk of malnutrition prepared by the Ministry of Health of Brazil^[23] shows that the implementation of ONT is able to reduce costs by R\$10,816.08 per death averted compared to enteral and parental nutritional therapies, while there are savings of R\$4,231.70 due to avoiding readmission and R\$283.06 for avoiding hospitalization.

Although doctors have the view that one of the benefits of ONS is the decrease in the chances of hospital readmission, most of the nursing team does not believe in this benefit. A study directed by Kaegi-Braun et al. (2021)^[24] found significantly lower rates of hospital readmission within 30 days in patients who received nutritional support; moreover, these patients had a lower mortality rate than those not receiving nutritional support.

In contrast to the results obtained by Brindisi et al. (2020)^[21] in which physicians classify food supplements as medication, there was unanimous agreement that ONS is a food supplement in the current study. Nevertheless, both groups claim that they did not have technical training regarding nutritional supplementation during their studies.

The nutritional risk criteria used in the questionnaires were based on a proprietary institutional tool. This study points out that several symptoms and conditions are neglected by professionals. For example, the presence of ascites, defined as the presence of free fluid of pathological origin in the abdominal cavity,^[25] was not often mentioned by the physicians or nursing staff as a criterion for patients to be at nutritional risk. This condition may be related to indigestion, as gastric capacity is reduced with an increased intra-abdominal pressure, which can lead to reduced food intake.^[26]

The professionals also did not believe that patients with comorbidities, such as diabetes mellitus, would benefit from ONS. According to the Guidelines of the Brazilian Society of Diabetes,^[27] oral supplements promote greater glycemic control by reducing

postprandial blood glucose and triglyceride levels, reducing insulin resistance, controlling the lipid profile, and reducing episodes of hypoglycemia. This finding indicates that further training of physicians and nurses regarding the benefits of ONS is needed.

Physicians participating in this study collectively believed that patients who present with weight loss are eligible to receive ONS; however, in clinical practice, most patients are referred for nutritional monitoring only when there is a severe loss of weight. There are also many patients who are not provided with nutritional consultation and do not receive a medical prescription for food support. A study by Lorton et al. (2020)^[28] has demonstrated that 40% of malnourished patients do not receive adequate nutritional support. According to the National Oncology Nutrition Consensus,^[29] oral supplements are great allies against weight loss, preventing it from occurring and, in some cases, even weight gain being noticed.

Although the literature is clear regarding the nutritional status depletion that systemic treatment can cause, physicians and nurses in this study believe that there are no such negative effects. However, Carniatio et al. (2018)^[30] have shown that patients undergoing concomitant treatment have a greater acute toxicity to the treatment. In a paper presented by Rebouças et al. (2011)^[31] radiotherapy combined with chemotherapy in patients diagnosed with head and neck cancer is linked to a higher frequency in grade 3 acute toxicity.

Physicians as well as nurses and technicians do not consider enteral and parenteral nutritional therapy at home as a risk factor for malnutrition; however, the presence of nutritional support is not synonymous with the treatment being adequate. Mazur et al. (2014)^[32] have reported that home enteral nutritional therapy has several factors that hamper the administration of tube feeding in the home context, for example, financial aspects can lead to the impossibility of purchasing formula, leading to an inadequate food supply and thus causing malnutrition. In addition, adaptation of family to a new daily life can also contribute to food insecurity.

Even without training on the proposed subject, most physicians and nursing staff notice the improvement in the nutritional status of patients who use ONS. However, even without adequate training, they felt able to guide patients regarding the consumption of ONS that they prescribed and administered, respectively. These findings corroborate the results presented by Spiro et al. (2016),^[33] who suggest that oncologists believe that the nutritional status of patients and nutritional interventions are relevant for patients undergoing anticancer treatment. Yet they are unable to identify patients at nutritional risk and to refer those who would benefit from nutritional therapy at an early stage. Their study also found that malnutrition is not widely recognized by other health professionals.

When nutritional care takes place in a multidisciplinary way, health outcomes are more favorable to the patient's quality of life. Such care should involve all health professionals such as dietitians, physicians, and nurses at all levels of care involving nutritional screening, assessment, planning, implementation, evaluation and ultimately the monitoring of evidence-based care delivery.^[34]

The study by Vrkić et al. (2022)^[35] points out that high-quality continuing medical education related to nutrition is capable of increasing medical knowledge and skills in relation to nutritional care, training professionals with greater confidence in the skills needed for nutritional interventions. This also mentions that the lack of training and nutritional education can have a vital impact on the nutritional information provided to patients, which may compromise the safety and efficiency of nutritional counseling, putting the patient at risk.

Continuing education in cancer nutrition is essential for every multidisciplinary team, since nutrition is an important part of treatment and not just a complement.^[36] Nutritional education interventions, such as the creation of workshops, protocols and discussion of clinical cases, increase the quality of nutritional therapy, this study also points out that there was an improvement in clinical practice after the application of training for physicians, impacting on a shorter period of fasting and greater achievement of nutritional goals.^[37]

Nutritional diagnosis must be carried out in parallel with disease diagnosis so that early nutritional intervention is possible, thus affording better clinical outcomes.^[38]

The present study has some limitations, such as the use of a questionnaire designed for this study, based on other similar research, but not validated. In addition, the research was conducted with only one medical specialty, since other specialties, such as surgeons, have greater knowledge about nutritional supplements due to institutional protocols that encompass this type of nutritional therapy. Finally, the nursing staff evaluated was only those who had care with the inpatient.

CONCLUSION

There is a great difference in knowledge about ONS among medical and nursing professionals. Interdisciplinary technical training is essential for the success of ONT as a tool for preventing and combating malnutrition. Although nutrition is not a part of the curriculum in other areas of health care, hospitals must carry out training to improve knowledge regarding this topic so that ONT can be implemented.

AUTHORS' CONTRIBUTIONS

TMM	Final approval of manuscript, Manuscript writing
CRK	Collection and assembly of data, Conception and design, Data analysis and interpretation, Final approval of manuscript, Manuscript writing
JPF	Final approval of manuscript, Manuscript writing

REFERENCES

- Toledo DO, Piovacari SMF, Horie LM, Matos LBN, Castro MG, Ceniccola GD, et al. Campanha 'Diga não à desnutrição': 11 passos importantes para combater a desnutrição hospitalar. *BRASPEN J.* 2018;33(1):86-100.
- Cederholm T, Bosaeus I, Barazzoni R, Bauer J, Van Gossum A, Klek S, et al. Diagnostic criteria for malnutrition. *ESPEN Consensus Statement. Clin Nutr.* 2015 Jun;34(3):335-40.
- Cederholm T, Jensen GL, Correia MITD, Gonzalez MC, Fukushima R, Higashiguchi T, et al. GLIM criteria for the diagnosis of malnutrition – a consensus report from the global clinical nutrition community. *Clin Nutr.* 2019 Feb;38(1):9-1.
- Uster A, Ruefenacht U, Ruelin M, Pless M, Siano M, Haefner M, et al. Influence of a nutritional intervention on dietary intake and quality of life in cancer patients: a randomized controlled trial. *Rev Nutr.* 2013 Nov/Dec;29(11-12):1342-9.
- Muscaritoli M, Lucia S, Farcomeni A, Lorusso V, Saracino V, Barone C, et al. Prevalence of malnutrition in patients at first medical oncology visit: the PreMiO study. *Oncotarget.* 2017 Aug;8(45):79884-96.
- Arends J, Bachmann P, Baracos V, Barthelemy N, Bertz H, Bozzetti F, et al. ESPEN guidelines on nutrition in cancer patients. *Clin Nutr.* 2017 Feb;36(1):11-48.
- Muscaritoli M, Molino A, Gioia G, Laviano A, Fanelli FR. The "parallel pathway": a novel nutritional and metabolic approach to cancer patients. *Intern Emerg Med.* 2011 Apr;6(2):105-12. DOI: <https://doi.org/10.1007/s11739-010-0426-1>
- Khalatbari-Soltani S, Marques-Vidal P. The economic cost of hospital malnutrition in Europe; a narrative review. *Clin Nutr ESPEN.* 2015 Jun;10(3):e89-e94.
- Zhang X, Tang T, Pang L, Sharma SV, Li R, Nyitray AG, et al. Malnutrition and overall survival in older adults with cancer: a systematic review and meta-analysis. *J Clin Oncol.* 2019 Nov;10(6):874-83.
- Segura A, Pardo J, Jara C, Zugazabeitia L, Carulla J, Peñas RDL, et al. An epidemiological evaluation of the prevalence of malnutrition in Spanish patients with locally advanced or metastatic cancer. *Clin Nutr.* 2005;24(5):801-814.
- Pelissaro E, Damo CC, Alves ALS, Calcing A, Kumpel DA. Avaliação do estado nutricional em pacientes idosos oncológicos internados em um hospital de alta complexidade do Norte do Rio Grande do Sul. *Sci Med.* 2016;26(2):ID22972.
- Cibulski TP, Becker T, Baldissera C, Basso T, Kumpel DA, et al. Adesão à terapia nutricional oral de pacientes com neoplasias de cabeça e pescoço. *BRASPEN J.* 2018;33(3):215-20.
- Piovacari SMF, Barrère APN. *Nutrição clínica na oncologia.* São Paulo: Atheneu; 2019.
- Silva MBR, Santos HMC, Oliveira AF. Características sensoriais e nutricionais de suplementos alimentares. *Ciênc Biol Saúde.* 2014;35:31-8.

15. Tanaka M, Marques APA, Costa VS, Piovacari SMF, Sandoval LCN. Terapia nutricional oral. In: Piovacari SMF, Toledo DO, Figueiredo EJA, eds. Equipe multiprofissional de terapia nutricional. São Paulo: Atheneu; 2017. p. 256-236.
16. Ministério da Saúde (BR). Agência Nacional de Vigilância Sanitária (Anvisa). Macrotema de Alimentos. 5ª ed. Brasília (DF): Ministério da Saúde/Anvisa; 2020.
17. Ravasco P. Nutrition in cancer patients. *J Clin Med*. 2019 Aug;8(8):1211.
18. Baldwin C, Spiro A, Ahern R, Emery PW. Oral nutritional interventions in malnourished patients with cancer: a systematic review and meta-analysis. *J Natl Cancer Inst*. 2012 Feb;104(5):371-85. DOI: <https://doi.org/10.1093/jnci/djr556>
19. Paccagnella A, Morello M, Mosto MC, Baruffi C, Marcon ML, Gava A, et al. Early nutritional intervention improves treatment tolerance and outcomes in head and neck cancer patients undergoing concurrent chemoradiotherapy. *Support Care Cancer*. 2010 Jul;18(7):837-45.
20. Philipson TJ, Snider JT, Lakdawalla DN, Stryckman B, Goldman DP. Impact of oral nutritional supplementation on hospital outcomes. *Am J Manag Care*. 2013 Feb;19(2):121-8.
21. Brindisi MC, Noacco A, Hansal AAB, Gentil CH. Delivery of oral nutrition supplement in hospital: Evaluation of professional practices in evaluation of nutritional status and representations of ONS by the caregivers and patients. *Clin Nutr ESPEN*. 2020 Feb;35:85-9.
22. Muscaritoli M, Molfino A, Gioia G, Laviano A, Fanelli FR. The "parallel pathway": a novel nutritional and metabolic approach to cancer patients. *Intern Emerg Med*. 2011;6(2):105-12.
23. Ministério da Saúde (BR). Secretaria de Ciência, Tecnologia, Inovação e Insumos Estratégicos em Saúde. Suplementação nutricional oral para pacientes clínicos ou cirúrgicos desnutridos ou em risco de desnutrição. Brasília (DF): Ministério da Saúde; 2021.
24. Kaegi-Braun N, Mueller M, Schuetz P, Mueller B, Kutz A. Evaluation of nutritional support and in-hospital mortality in patients with malnutrition. *JAMA Network Open*. 2021;4(1):e2033433.
25. Júnior DRA, Galvão FHF, Santos SA. Ascite – Estado baseado em evidências. *Rev Assoc Med Bras*. 2009;55(4):489-96.
26. Gregorini, FG, Stanich P, Freitas MMT. Avaliação do estado nutricional em pacientes hospitalizados com cirrose hepática. *BRASPEN J*. 2016;31(4):299-304.
27. Golbert A, Vasques ACJ, Faria ACRA, Lottenberg AMP, Joaquim AG, Vianna AGD. Diretrizes sociedade brasileira de diabetes. São Paulo (SP): Clannad; 2019.
28. Lorton CM, Griffin O, Higgins K, Roulston F, Stewart G, Gough N, et al. Late referral of cancer patients with malnutrition to dietitians: a prospective study of clinical practice. *Support Care Cancer*. 2020 May;28(5):2351-60.
29. Ministério da Saúde (BR). Instituto Nacional de Câncer José Alencar Gomes da Silva (INCA). Consenso nacional de nutrição oncológica. 2ª ed. Rio de Janeiro (RJ): INCA; 2016
30. Carniatio LN, Miola TM, Chulam TC. Evolution of nutritional status of patients with head and neck cancer during radiotherapy or radiotherapy concerning chemotherapy. *BRASPEN J*. 2018;33(2):141-6.
31. Rebouças LM, Callegaro E, Gil GOB, Silva MLG, Maia MAC, Salvajoli JV. Impacto da nutrição enteral na toxicidade aguda e na continuidade do tratamento dos pacientes com tumores de cabeça e pescoço submetidos a radioterapia com intensidade modulada. *Radiol Bras*. 2011 Jan/Feb;44(1):42-6.
32. Mazur CE, Schmidt ST, Rigon AS, Schieferdecker MEM. Terapia nutricional enteral domiciliar: interface entre direito humano à alimentação adequada e segurança alimentar e nutricional. *Demetra*. 2014;9(3):757-69.
33. Spiro A, Baldwin C, Patterson A, Thomas J, Andreyev HJN. The views and practice of oncologists towards nutritional support in patients receiving chemotherapy. *Br J Cancer*. 2016 Aug;95(4):431-4.
34. Xiaoyue X, Parker D, Ferguson C, Hickman L. Where is the nurse in nutritional care?. *Contemp Nurse*. 2017 Jun;53(3):267-70.
35. Vrkić A, Grujić M, Jovičić-Bata J, Novaković B. Nutritional knowledge, confidence, attitudes toward nutritional care and nutrition counseling practice among general practitioners. *Healthcare*. 2022 Sep;10(11):2222.
36. Liposits G, Orreval Y, Kaasa S, Osterlund P, Ceder T. Nutrition in cancer care: a brief, practical guide with a focus on clinical practice. *J Oncol Pract*. 2021;17(7):992-8.
37. Ferreira HCC, Rodrigues PL. Avaliação do conhecimento médico sobre terapia nutricional parenteral em um hospital público do Distrito Federal. *BRASPEN J*. 2017;32(4):387-93.
38. Muscaritoli M, Arends J, Bachmann P, Baracos V, Barthelemy N, Bertz H, et al. ESPEN practical guideline: clinical nutrition in cancer. *Clin Nutr*. 2021;40:2913-98.

Appendix 1 - Questionnaire for Nursing Staff

Name: _____

 Nurse Nursing assistant Nursing technician

<p>1. Do you know what an oral nutritional supplement (ONS) is?</p> <input type="checkbox"/> Medication <input type="checkbox"/> Meal replacement <input type="checkbox"/> Meal complement <input type="checkbox"/> I do not know <input type="checkbox"/> Others: _____
<p>2. Was this topic addressed during your professional training?</p> <input type="checkbox"/> Yes <input type="checkbox"/> No <input type="checkbox"/> I do not know
<p>3. Who is this supplement for?</p> <input type="checkbox"/> Underweight patients <input type="checkbox"/> Patients with associated diseases <input type="checkbox"/> I do not know <input type="checkbox"/> Malnourished patients <input type="checkbox"/> All cancer patients <input type="checkbox"/> Others: _____
<p>4. What criteria would you use to classify a patient at nutritional risk?</p> <input type="checkbox"/> Weight loss <input type="checkbox"/> Low food acceptance <input type="checkbox"/> Dysphagia <input type="checkbox"/> Diarrhea <input type="checkbox"/> Vomiting <input type="checkbox"/> Mucositis <input type="checkbox"/> Head and neck cancer <input type="checkbox"/> Ascites <input type="checkbox"/> Enteral/Parenteral Home Therapy <input type="checkbox"/> Chemotherapy concomitant with radiotherapy <input type="checkbox"/> Others: _____
<p>5. What benefits do you believe ONS brings to cancer patients?</p> <input type="checkbox"/> Weight gain <input type="checkbox"/> Better food intake <input type="checkbox"/> Better tolerance to treatment <input type="checkbox"/> Lower chances of hospital readmission <input type="checkbox"/> None <input type="checkbox"/> Others: _____
<p>6. Do you notice an improvement in the patient's general condition and/or food acceptance while using the ONS? (e.g., weight gain)</p> <input type="checkbox"/> Yes <input type="checkbox"/> No <input type="checkbox"/> I do not know how to answer
<p>7. By whom is ONS prescribed?</p> <input type="checkbox"/> Oncologist <input type="checkbox"/> Nutritionist <input type="checkbox"/> I do not know <input type="checkbox"/> Nutritional Doctor <input type="checkbox"/> Others: _____
<p>8. Who is the professional responsible for delivering the ONS in your department? (Can check more than one option)</p> <input type="checkbox"/> Nurse <input type="checkbox"/> Nursing technician <input type="checkbox"/> Nursing assistant <input type="checkbox"/> Others: _____
<p>9. When administering the ONS to the patient, do you guide him/her? (Indicate if you are responsible for the delivery)</p> <input type="checkbox"/> Yes <input type="checkbox"/> No <input type="checkbox"/> I do not know
<p>10. If not, why?</p> <input type="checkbox"/> It is another professional's responsibility <input type="checkbox"/> I do not know how to assess the need for the supplement <input type="checkbox"/> The nutritional status of the patient is not relevant to his/her treatment <input type="checkbox"/> Others: _____
<p>11. What do you do if the patient refuses the ONS?</p> <input type="checkbox"/> I advise him/her to talk to the doctor <input type="checkbox"/> I advise him/her to talk to the nutritionist <input type="checkbox"/> I communicate with the doctor <input type="checkbox"/> I communicate with the nutritionist <input type="checkbox"/> Others: _____
<p>Personal comments:</p>

Appendix 2 - Questionnaire for Medical Staff

Name: _____

 Senior doctor Resident doctor

<p>1. Do you know what oral nutritional supplement (ONS) is?</p> <p><input type="checkbox"/> Medication <input type="checkbox"/> Meal replacement <input type="checkbox"/> Meal complement</p> <p><input type="checkbox"/> I do not know <input type="checkbox"/> Others: _____</p>
<p>2. Was this topic addressed during your professional training?</p> <p><input type="checkbox"/> Yes <input type="checkbox"/> No <input type="checkbox"/> I do not know</p>
<p>3. Do you prescribe or have you already prescribed an ONS?</p> <p><input type="checkbox"/> Yes <input type="checkbox"/> No</p>
<p>4. Based on which diagnosis do you prescribe an ONS?</p> <p><input type="checkbox"/> Underweight patients <input type="checkbox"/> Patients with associated diseases <input type="checkbox"/> I do not know</p> <p><input type="checkbox"/> Malnourished patients <input type="checkbox"/> All cancer patients <input type="checkbox"/> Others: _____</p>
<p>5. What criteria would you use to classify a patient at nutritional risk?</p> <p><input type="checkbox"/> Weight loss <input type="checkbox"/> Low food acceptance <input type="checkbox"/> Dysphagia <input type="checkbox"/> Diarrhea <input type="checkbox"/> Vomiting</p> <p><input type="checkbox"/> Mucositis <input type="checkbox"/> Head and neck cancer <input type="checkbox"/> Ascites <input type="checkbox"/> Enteral/Parenteral Home Therapy</p> <p><input type="checkbox"/> Chemotherapy concomitant with radiotherapy <input type="checkbox"/> Others: _____</p>
<p>5. What benefits do you believe the ONS brings to cancer patients?</p> <p><input type="checkbox"/> Weight gain <input type="checkbox"/> Better food intake <input type="checkbox"/> Better tolerance to treatment</p> <p><input type="checkbox"/> Lower chances of hospital readmission <input type="checkbox"/> None <input type="checkbox"/> Others: _____</p>
<p>6. When prescribing an ONS for a patient, do you guide him/her?</p> <p><input type="checkbox"/> Yes <input type="checkbox"/> No <input type="checkbox"/> I do not prescribe ONS</p>
<p>7. If not, why?</p> <p><input type="checkbox"/> It is another professional's responsibility <input type="checkbox"/> I do not know how to assess the need for the supplement</p> <p><input type="checkbox"/> The nutritional status of the patient is not relevant to his/her treatment <input type="checkbox"/> Others: _____</p>
<p>8. What benefits do you believe the ONS brings to cancer patients?</p> <p><input type="checkbox"/> Weight gain <input type="checkbox"/> Better food intake <input type="checkbox"/> Better tolerance to treatment</p> <p><input type="checkbox"/> Lower chances of hospital readmission <input type="checkbox"/> None <input type="checkbox"/> Others: _____</p>
<p>9. Do you notice an improvement in the patient's general condition and/or food acceptance while using the ONS? (e.g., weight gain)</p> <p><input type="checkbox"/> Yes <input type="checkbox"/> No <input type="checkbox"/> I do not know how to answer</p>
<p>10. What do you do if the patient refuses the ONS?</p> <p><input type="checkbox"/> I suspend treatment <input type="checkbox"/> I advise him/her to talk to the nutritionist <input type="checkbox"/> I communicate with the nutritionist</p> <p><input type="checkbox"/> I reinforce the importance of the ONS <input type="checkbox"/> Others: _____</p>
<p>Personal comments:</p>